arXiv:1409.5367v1 [math.NT] 18 Sep 2014

DIFFERENTIAL MODULAR FORMS ATTACHED TO NEWFORMS MOD p

ALEXANDRU BUIUM

Abstract. In a previous paper [7] we attached to classical complex newforms f of weight 2 certain δ_p-modular forms $f^\sharp$ (in the sense of [4, 6]) of order 2 and weight 0; the forms $f^\sharp$ can be viewed as "dual" to f and played a key role in some of the applications of the theory [8]. The aim of this paper is to provide a higher weight version of this "$\sharp$-duality," by attaching to classical newforms mod p, $\overline{f}$, of weight κ between 3 and p, δ_π-modular forms $f^\sharp$ of order 2 and weight $-\kappa'$, with κ' between 1 and $p-2$.

1. Introduction and main result

In a series of papers beginning with [3] an arithmetic analogue of differential equations was developed in which derivatives of functions were replaced by Fermat quotient operators δ_π; the latter act on complete discrete valuation rings R_π of unequal characteristic and uniformizer π, by the formula $\delta_\pi x := \frac{\phi(x)-x^p}{\pi}$, where $\phi : R_\pi \to R_\pi$ is a ring endomorphism whose reduction mod π is the p-power Frobenius. In particular spaces of δ_π-modular forms were introduced and studied (cf. [4, 6] for $\pi = p$ and [10] for arbitrary π).

A basic construction in [7] attaches to classical newforms $f = \sum a_n q^n$ over $\mathbb{C}$, of level N (i.e. on $\Gamma_1(N)$) and weight 2, some δ_p-modular forms $f^\sharp$ of level N, order 2, and weight 0, with δ_p-Fourier expansion nicely expressible in terms of $f^{(-1)}(q) := \sum \frac{a_n}{n} q^n$. The forms $f^\sharp$ in [7] are arithmetic-differential objects that have no classical analogue but, rather, can be viewed as "dual" to the classical objects f; by the way, the forms $f^\sharp$ introduced in [7] played a key role in [8] where the theory of arithmetic differential equations was used to prove finiteness results for Heegner-like points.

The aim of this paper is to extend this "$\sharp$-duality" to the higher weight case. Specifically let N be an integer with $(N, p) = 1, N > 4$, let k be an algebraic closure of $\mathbb{F}_p$, and let $\overline{f}$ be a newform of level N, over k, of weight κ, with $3 \leq \kappa \leq p$. Then, by a construction due to Serre [14], there exists a newform f of level Np over $\mathbb{C}$, of weight 2, lifting $\overline{f}$, whose character is compatible, in a sense to be reviewed later, with the Teichmüller character. (The compatibility requires an embedding of the field K_f generated by the Fourier coefficients of f into some $K_\pi := R_\pi[1/p]$; the considerations below will be valid for a suitable embedding of the Galois closure of K_f into a suitable K_π.) Let us say that f is a *Serre lift* of $\overline{f}$. Also let us say that an integer κ' is a *conjugate* of an integer $3 \leq \kappa \leq p$ if $1 \leq \kappa' \leq p-2$ and there exists an integer c coprime to $p-1$ such that $\kappa' \equiv c(2-\kappa) \bmod p-1$. We recall from [6, 10] that if $q, \delta_\pi q, ..., \delta_\pi^n q$ are variables then any δ_π-modular form g possesses a δ_π-Fourier expansion $E(g)$ in $R_\pi((q))[\delta_\pi q, ..., \delta_\pi^n q]^\wedge$ (where $\wedge$ always means here p-adic

2000 *Mathematics Subject Classification.* 11E95, 11F37.

completion). On the other hand ϕ acts on $K_\pi[[q]]^\infty := \cup_n K_\pi[[q, \delta_\pi q, ..., \delta_\pi^n q]]$ by $\phi(q) = q^p + \pi\delta_\pi q$, $\phi(\delta_\pi q) = (\delta_\pi q)^p + \pi\delta_\pi^2 q$, etc. In particular $K_\pi[[q]]^\infty$ has a natural structure of left module over the polynomial ring $K_\pi[\phi]$. So in particular, for any polynomial $P(\phi) \in K_\pi[\phi]$ and any series $g(q) \in K_\pi[[q]]$ (e.g., for $g = f^{(-1)}(q)$) it makes sense to consider the series $P(\phi)g(q) \in K_\pi[[q]]^\infty$. The above concepts will be reviewed in detail in the body of the paper. We will then prove the following:

Theorem 1.1. *For any newform $\overline{f}$ over k, of level N and weight κ, with $3 \leq \kappa \leq p$, there exists a non-zero δ_π-modular form $f^\sharp$ of level N, order 2, and weight $-\kappa'$, with κ' a conjugate of κ, such that $f^\sharp$ has a δ_π-Fourier expansion of the form*

$$E(f^\sharp) = \sum_\sigma P_\sigma(\phi)(f^{(-1)}(q))^\sigma,$$

with $P_\sigma(\phi)$ polynomials in $K_\pi[\phi]$, f a Serre lift of $\overline{f}$, and σ running through the set of all embeddings of K_f into $\mathbb{C}$.

The strategy of proof of our Theorem is as follows. The Serre lift f being of weight 2 and level Np one can use the method in [7] to attach to it an element $f^\sharp$ in the ring of *δ_π-modular forms of level Np and weight* 0. Using a construction from [10] one can interpret this $f^\sharp$ as a sum of δ_π-modular forms of level N and weights $0, -1, -2, ..., -p+2$. Finally, an analysis of the action of the diamond operators shows that this sum of δ_π-modular forms reduces to one term only; that term turns out to have weight $-\kappa'$ with κ' conjugate to κ. Note that each given κ has a priori several conjugates and the theorem does not tell which of these conjugates gives the weight of $f^\sharp$; be that as it may the weight $-\kappa'$ of $f^\sharp$ is always non-zero (in contrast to the weight of $f^\sharp$ in [7, 10] which is equal to 0). Finally note that, as in [7], $f^\sharp$ in our Theorem 1.1 is not unique.

The plan of the paper is as follows. In section 2 we review π-jet spaces and δ_π-modular forms following [3, 10]. In section 3 we review (and complement) some (complex and arithmetic) facts about classical modular forms following [14, 10]. In section 4 we construct our forms $f^\sharp$. In section 5 we use some of the tools developed in the paper to construct a remarkable homomorphism. Indeed, we will recall, in that section, the definition of the ring $S_\heartsuit^\infty$ of *Igusa δ_π-modular functions of level N* introduced in [9] and we will prove:

Theorem 1.2. *There is a natural homomorphism from the ring of δ_π-modular forms of level Np and weight* 0 *to the ring $S_\heartsuit^\infty \otimes_{R_p} R_\pi$. This homomorphism commutes with δ_π and raises orders by* 1.

Cf. Theorem 5.1 for a more precise statement.

Acknowledgement. The author would like to acknowledge partial support from the Simons Foundation (award 311773), from the Institut des Hautes Etudes Scientifiques in Bures sur Yvette, and from the Romanian National Authority for Scientific Research (CNCS - UEFISCDI, project number PN-II-ID-PCE-2012-4-0201).

2. Review of differential modular forms

In this section we review the basic definitions of π-jet spaces and δ_π-modular forms introduced in [3, 4, 10]. Throughout this paper $p \geq 5$ is a fixed prime. For any

ring A we denote the p-adic completion of A by $\widehat{A}$. We denote by $R_p = \widehat{\mathbb{Z}_p^{ur}}$ the p-adic completion of the maximum unramified extension of $\mathbb{Z}_p$. We set $K_p = R_p[1/p]$ (fraction field of R_p) and $k = R_p/pR_p$ (residue field of R_p); so k is an algebraic closure of $\mathbb{F}_p$.

We need a ramified version of the above (which is slightly more general than the one encountered in [3, 10]). Indeed let $F/\mathbb{Q}$ be a normal finite extension with ring of integers $\mathcal{O}_F$ and let $\wp \subset \mathcal{O}_F$ be a prime dividing p. There exists a (not necessarily unique) ring automorphism $\phi = \phi_\wp$ of $\mathcal{O}_F$ preserving $\wp$ and inducing the p-power Frobenius on $\mathcal{O}_F/\wp$. Let $\mathcal{O}_\wp$ be the completion of the localization of $\mathcal{O}$ at $\wp$ and $\pi \in \mathcal{O}_F$ a uniformizer of $\mathcal{O}_\wp$. Then $\phi : \mathcal{O}_F \to \mathcal{O}_F$ induces an automorphism of $\mathcal{O}_\wp$ whose reduction mod π is the p-power Frobenius on $\mathcal{O}_\wp/(\pi)$. For any discrete valuation ring $\mathcal{O}'$ which is a finite unramified extension of $\mathcal{O}_\wp$, $\mathcal{O}'$ is étale over $\mathcal{O}_\wp$ so $\phi : \mathcal{O}_\wp \to \mathcal{O}_\wp$ lifts uniquely to a ring automorphism $\phi : \mathcal{O}' \to \mathcal{O}'$ whose reduction mod π is the p-power Frobenius. Taking limit and completion we get an automorphism ϕ of the ring $\widehat{\mathcal{O}_\wp^{ur}}$, completion of the maximum unramified extension of $\mathcal{O}_\wp$, whose reduction mod π is the p-power Frobenius on the residue field. Let $K_\pi = K_p(\pi)$ and let R_π be the valuation ring of K_π. Since $R_\pi \subset \widehat{\mathcal{O}_\wp^{ur}}$ is an unramified extension of complete discrete valuation rings with the same residue field it follows that $R_\pi = \widehat{\mathcal{O}_\wp^{ur}}$ so ϕ is an autmorphism of R_π. We let e be the ramification index of $R_p \subset R_\pi$. Note that the resulting embedding $\rho : \mathcal{O}_F \to R_\pi$ depends on F and $\wp$ only.

Consider the polynomial

$$C_\pi(X,Y) := \pi^{-1}(X^p + Y^p - (X+Y)^p) \in \mathbb{Z}[p/\pi][X,Y].$$

A π-*derivation* from an $\mathbb{Z}[p/\pi]$-algebra A into an $A-$algebra B is a map $\delta_\pi : A \to B$ such that $\delta_\pi(1) = 0$ and

$$\begin{aligned} \delta_\pi(x+y) &= \delta_\pi x + \delta_\pi y + C_\pi(x,y) \\ \delta_\pi(xy) &= x^p \cdot \delta_\pi y + y^p \cdot \delta_\pi x + \pi \cdot \delta_\pi x \cdot \delta_\pi y, \end{aligned}$$

for all $x, y \in A$. For a $\pi-$derivation we denote by $\phi : A \to B$ the map $\phi(x) = x^p + \pi\delta_\pi x$; then ϕ is a ring homomorphism. A δ_π-*prolongation sequence* is a sequence $S^* = (S^n)_{n\geq 0}$ of $\mathbb{Z}[p/\pi]$-algebras S^n, $n \geq 0$, together with $\mathbb{Z}[p/\pi]$-algebra homomorphisms $\varphi : S^n \to S^{n+1}$ and $\pi-$derivations $\delta_\pi : S^n \to S^{n+1}$ (where S^{n+1} is viewed as an S^n-algebra via φ) such that $\delta_\pi \circ \varphi = \varphi \circ \delta_\pi$ on S^n for all n. A morphism of δ_π-prolongation sequences, $u^* : S^* \to \tilde{S}^*$ is a sequence $u^n : S^n \to \tilde{S}^n$ of $\mathbb{Z}[p/\pi]$-algebra homomorphisms such that $\delta_\pi \circ u^n = u^{n+1} \circ \delta_\pi$ and $\varphi \circ u^n = u^{n+1} \circ \varphi$. Let W be the ring of polynomials $\mathbb{Z}[\phi]$ in the indeterminate ϕ. For $w = \sum_{i=0}^r a_i \phi^i \in W$ (respectively for w with $a_i \geq 0$), S^* a δ_π-prolongation sequence, and $x \in (S^0)^\times$ (respectively $x \in S^0$) we can consider the element $x^w := \prod_{i=0}^r \varphi^{r-i}\phi^i(x)^{a_i} \in (S^r)^\times$ (respectively $x^w \in S^r$). On the other hand we may consider the $\pi-$derivation $\delta_\pi : R_\pi \to R_\pi$ given by $\delta_\pi x = (\phi(x) - x^p)/\pi$. One can consider the δ_π-prolongation sequence R_π^* where $R_\pi^n = R_\pi$ for all n. By a δ_π-*prolongation sequence over* R_π we understand a prolongation sequence S^* equipped with a morphism $R_\pi^* \to S^*$. From now on all our δ_π-prolongation sequences are assumed to be over R_π. For any affine R_π-scheme of finite type $Y = Spec\ A$ there exists a (unique) δ_π-prolongation sequence, $A^* = (A^n)_{n\geq 0}$, with $A^0 = A$ such that for any δ_π-prolongation sequence B^* and any R_π-algebra homomorphism $u : A \to B^0$ there exists a unique morphism of δ_π-prolongation sequences $u^* : A^* \to B^*$ with $u^0 = u$. We define the π-*jet spaces*

$J^n_\pi(Y)$ of Y as the formal schemes $J^n_\pi(Y) := Spf\ \hat{A}^n$. This construction globalizes to the case when Y is not necessarily affine (such that the construction commutes, in the obvious sense, with open immersions). A δ_π-*function* on Y (of order n) is an element $f \in \mathcal{O}(J^n_\pi(Y))$, equivalently a morphism $f : J^n_\pi(Y) \to \widehat{\mathbb{A}^1}$ over R_π; such a map induces a map $f_* : Y(R_\pi) \to R_\pi$. (Indeed any R_π-point of Y canonically lifts, by the universality property, to an R_π-point of $J^n_\pi(Y)$ which can then be mapped by f_* to an R_π-point of the affine line.) If Y is smooth f_* uniquely determines f; we then write $f_* = f$ and we also refer to f_* as a δ_π-*function*. If $Y = G$ is a smooth group scheme over R_π then a δ_π-*character* of G is a homomorphism $f : J^n_\pi(G) \to \widehat{\mathbb{G}_a}$; such an f induces a group homomorphism $f = f_* : G(R_\pi) \to R_\pi$, still referred to as a δ_π-character. Note that the π-jet spaces $J^n_\pi(Y)$ only depend on the π-adic completion of Y and not on Y so one can introduce π-jet spaces $J^n_\pi(\mathcal{Y})$ attached to formal π-adic schemes $\mathcal{Y}$ over R_π which are locally π-adic completions of schemes of finite type over R_π.

Next we review differential modular forms (δ_p-modular forms and δ_π-modular forms) following [4, 10]. Recall the modular curve $X_1(N)_{R_p}$ over R_p with $(N, p) = 1$, $N > 4$; it is a smooth curve and it carries a line bundle L such that the spaces of sections of L^κ identify with the spaces of modular forms on $\Gamma_1(N)$ defined over R_p of weight κ; cf. [14], p. 450; in loc. cit. L was denoted by ω. The curve $X_1(N)_{R_p}$ contains two (disjoint) closed subsets of interest: the *cusp locus* (*cusps*) and the *supersingular locus* (*ss*). On $Y_1(N) = X_1(N)\backslash(cusps)$ the line bundle L identifies with $u_*\Omega^1_{E/Y_1(N)}$ where $u : E \to Y_1(N)$ is the corresponding universal elliptic curve. Next consider an affine open set $X \subset X_1(N)_{R_p}$, disjoint from the supersingular locus (but not necessarily from the cusps!), and consider the restriction of L to X which we continue to denote by L. We can consider the affine X-scheme $V := Spec\left(\bigoplus_{n\in\mathbb{Z}} L^{\otimes n}\right) \to X$; then V is naturally a principal (Zariski locally trivial) $\mathbb{G}_m$-bundle. A δ_π-*modular function* on X, of level N and order r is a δ_π-function $V(R_\pi) \to R_\pi$ of order r. We say that a δ_π-modular function f has *weight* $w \in W$ if $\lambda \star f = \lambda^w f$ for $\lambda \in R^\times_\pi$ where $\star$ is the natural $\mathbb{G}_m$-action on V; δ_π-modular functions possessing weights are called δ_π-*modular forms*. We denote by $M^r_\pi(w)$ the spaces of δ_π-modular forms of weight w and order r. Recall from [10] that, for variables denoted by $q, \delta_\pi q, ..., \delta^r_\pi q$, one has natural injective δ_π-Fourier expansion maps

$$E : M^r(w) \to R_\pi((q))^r := R_\pi((q))[\delta_\pi q, ..., \delta^r_\pi q]\hat{},$$

induced by the universality property of $(\mathcal{O}(J^r_\pi(V)))_{r\geq 0}$ and the obvious structure of prolongation sequence of $(R_\pi((q))^r)_{r\geq 0}$.

3. Review of classical modular forms

In this section we review (and complement) some basic facts about classical modular forms in [12, 14, 10].

3.1. **Complex modular forms** [12]**.** We denote by $M_m(\Gamma_1(M), \mathbb{C})$ the space of classical complex modular forms of weight m on $\Gamma_1(M)$ and by $T_m(n)$ and $\langle n\rangle$, $n \geq 1$, the Hecke operators and the diamond operators acting on this space. The diamond operators belong to the ring generated by the Hecke operators. An eigenform is an element $f \in M_m(\Gamma_1(M), \mathbb{C})$ which is an eigenvector for all $T_m(n)$ (and hence also for all $\langle n\rangle$). Let $S_m(\Gamma_1(M), \mathbb{C})$ be the space of cusp forms. An eigenform has $a_1 \neq 0$. If a cuspidal eigenform $f = \sum a_n q^n$ is normalized by $a_1 = 1$ then

$T_m(n)f = a_n f$ for $n \geq 1$ and there exists a Dirichlet character ϵ mod M such that $\langle n \rangle f = \epsilon(n) f$ for $n \geq 1$. Let f be a normalized cuspidal eigenform as above; one lets $\mathcal{O}_f \subset \mathbb{C}$ denote the ring generated by all a_n (it is finitely generated) and one lets K_f be the fraction field of $\mathcal{O}_f$. The Dirichlet character ϵ attached to f takes then values in $\mathcal{O}_f$. A newform is a cuspidal normalized eigenform such that the system $\{a_l; l \not| M\}$ does not occur for a cuspidal normalized eigenform form of the same weight and smaller level dividing M. Let $X_1(M)$ be the complete complex modular curve for $\Gamma_1(M)$; as an algebraic curve it is defined over $\mathbb{Q}$ hence its Jacobian $J_1(M)$ is also defined over $\mathbb{Q}$. The diamond operators act on $X_1(M)$ and are defined over $\mathbb{Q}$ and hence induce endomorphisms of $J_1(M)$ defined over $\mathbb{Q}$. We recall the following basic:

Theorem 3.1. *(Eichler-Shimura) Let $f \in S_2(\Gamma_1(M), \mathbb{C})$ be a newform with character ϵ. Then there exists an abelian variety $A_{f/\mathbb{Q}}$ over $\mathbb{Q}$, a ring homomorphism*

$$\iota : \mathcal{O}_f \to End(A_{f/\mathbb{Q}}/\mathbb{Q}),$$

and a dominant homomorphism

$$\pi : J_1(M)_{\mathbb{Q}} \to A_{f/\mathbb{Q}},$$

defined over $\mathbb{Q}$, such that for any prime l we have commutative diagrams

$$\begin{array}{ccc} J_1(M)_{\mathbb{Q}} & \stackrel{T(l)_*}{\longrightarrow} & J_1(M)_{\mathbb{Q}} \\ \pi \downarrow & & \downarrow \pi \\ A_{f/\mathbb{Q}} & \stackrel{\iota(a_l)}{\longrightarrow} & A_{f/\mathbb{Q}} \end{array} \qquad \begin{array}{ccc} J_1(M)_{\mathbb{Q}} & \stackrel{\langle l \rangle_*}{\longrightarrow} & J_1(M)_{\mathbb{Q}} \\ \pi \downarrow & & \downarrow \pi \\ A_{f/\mathbb{Q}} & \stackrel{\iota(\epsilon(l))}{\longrightarrow} & A_{f/\mathbb{Q}} \end{array}$$

Moreover the image of the pull-back map

$$\pi^* : H^0(A_{f/\mathbf{C}}, \Omega) \to H^0(J_1(N)_{\mathbf{C}}, \Omega) \simeq S_2(\Gamma_1(N), \mathbf{C})$$

is the $\mathbf{C}-$linear span of $\{f^\sigma \mid \sigma : K_f \to \mathbf{C}\}$.

Here $T(l)_*, \langle l \rangle_* \in End(J_1(M)_{\mathbb{Q}}/\mathbb{Q})$ are the naturally induced Hecke and diamond operators.

3.2. **Model of $X_1(Np)$.** For modular curves and forms we use the standard notation from [14]. Let $N > 4$, $(N, p) = 1$. Recall that the modular curve $X_1(Np)$ over $\mathbb{C}$ has a model (still denoted by $X_1(Np)$ in what follows) over $\mathbb{Z}[1/N, \zeta_p]$ considered in [14], p. 470. Recall some of the main properties of $X_1(Np)$. First $X_1(Np)$ is a regular scheme proper and flat of relative dimension 1 over $\mathbb{Z}[1/N, \zeta_p]$ (where ζ_p is a primitive p-th root of unity) and is smooth over $\mathbb{Z}[1/Np, \zeta_p]$. Also the special fiber of $X_1(Np)$ at p is a union of two smooth projective curves I and I' crossing transversally at a finite set Σ of points. Furthermore I is isomorphic to the Igusa curve $I_1(N)$ in [14], p. 160, so I is the smooth compactification of the curve classifying triples (E, α, β) with E an elliptic curve over a scheme of characteristic p, and $\alpha : \mu_N \to E$, $\beta : \mu_p \to E$ are embeddings (of group schemes). Similarly I' is the smooth compactification of the curve classifying triples (E, α, b) with E an elliptic curve over a scheme of characteristic p, and $\alpha : \mu_N \to E$, $b : \mathbb{Z}/p\mathbb{Z} \to E$ are embeddings. Finally Σ corresponds to the supersingular locus on the corresponding curves.

Let now $F_0 = \mathbb{Q}(\zeta_p)$ and $\pi_0 = 1 - \zeta_p$ and consider the embedding of $\mathbb{Z}[\zeta_N, \zeta_p, 1/N]$ into R_{π_0} (hence of $\mathbb{Z}[\zeta_N, 1/N]$ into R_p). Denote by $X_1(Np)_{R_{\pi_0}}$ the base change of

$X_1(Np)$ to R_{π_0}. Recall that the modular curve $X_1(N)$ over $\mathbb{C}$ has a natural smooth projective model (still denoted by $X_1(N)$) over $\mathbb{Z}[1/N]$ such that

$$Y_1(N) := X_1(N)\backslash(cusps)$$

parameterizes pairs (E, α) consisting of elliptic curves E with an embedding $\alpha : \mu_N \to E$. The morphism $X_1(Np) \to X_1(N)$ over $\mathbb{C}$ induces a morphism

$$e : X_1(Np)_{R_{\pi_0}}\backslash\Sigma \to X_1(N)_{R_{\pi_0}}\backslash(ss)$$

over R_{π_0}, where (ss) is the supersingular locus in the closed fiber of $X_1(N)_{R_{\pi_0}}$. (See [10] for details.) Let $X \subset X_1(N)_{R_p}\backslash(ss)$ be an affine open set, $X_{R_{\pi_0}} := X \otimes_{R_p} R_{\pi_0} \subset X_1(N)_{R_{\pi_0}}\backslash(ss)$ its base change to R_{π_0}, and $X_! := e^{-1}(X_{R_{\pi_0}})$. Denote by $\mathcal{X}_{R_{\pi_0}}$ the π_0-adic completion of $X_{R_{\pi_0}}$. Also note that the π_0-adic completion of $X_!$ has two connected components; let $\mathcal{X}_!$ be the component whose reduction mod π_0 is contained in $I\backslash\Sigma$. We get a morphism $e : \mathcal{X}_! \to \mathcal{X}_{R_{\pi_0}}$.

Now the diamond operators $\langle n\rangle$ on $X_1(Np)_{\mathbb{Q}}$ give rise to diamond operators $\langle n\rangle_N$ (for $n \in (\mathbb{Z}/N\mathbb{Z})^\times$) and $\langle d\rangle_p$ (for $d \in (\mathbb{Z}/p\mathbb{Z})^\times$) on $X_1(Np)_{\mathbb{Q}}$ corresponding to the Chinese remainder theorem spitting $(\mathbb{Z}/Np\mathbb{Z})^\times \simeq (\mathbb{Z}/N\mathbb{Z})^\times \times (\mathbb{Z}/p\mathbb{Z})^\times$. Recall from [14] that $G := (\mathbb{Z}/p\mathbb{Z})^\times$ acts via $\langle\ \rangle_p$ on the covering $X_1(Np) \to X_1(N)$ over $\mathbb{Z}[1/N, \zeta_p]$; this action preserves the Igusa curve I and induces on I the usual diamond operators. So G acts on the covering $e : \mathcal{X}_! \to \mathcal{X}_{R_{\pi_0}}$. Also, as a matter of notation, if $\epsilon : (\mathbb{Z}/Np\mathbb{Z})^\times \to \mathbb{C}$ is a Dirichlet character we denote by $\epsilon_N : (\mathbb{Z}/N\mathbb{Z})^\times \to \mathbb{C}$ and $\epsilon_p : (\mathbb{Z}/p\mathbb{Z})^\times \to \mathbb{C}$ the induced Dirichlet characters.

3.3. **Model of $J_1(Np)$.** In the discussion below we continue to assume that $\pi_0 = 1-\zeta_p$ and we continue to consider the embedding embedding $\mathbb{Z}[\zeta_N, \zeta_p, 1/N] \to R_{\pi_0}$. Let $A_{K_{\pi_0}} = J_1(Np)_{K_{\pi_0}}$ be the Jacobian of $X_1(Np)_{K_{\pi_0}}$ over K_{π_0}. Let $A_{R_{\pi_0}}$ be the Néron model of $A_{K_{\pi_0}}$ over R_{π_0}; cf. [2], p. 12. By the Néron property the Abel-Jacobi map $X_1(Np)_{R_{\pi_0}} \to J_1(Np)_{R_{\pi_0}}$ that sends ∞ into 0 can be extended to a morphism

$$X_1(Np)_{R_{\pi_0}}\backslash\Sigma \to A_{R_{\pi_0}}$$

and hence induces a morphism

$$\mathcal{X}_! \to \mathcal{A} := (A^0_{R_{\pi_0}})^\wedge$$

where $A^0_{R_{\pi_0}}$ is the connected component of $A_{R_{\pi_0}}$. Note that, since $\pi_0 = 1 - \zeta_p$, one has by [2], pp. 246 and 267, that $A^0_{R_{\pi_0}}$ coincides with $Pic^0_{X_1(Np)/R_{\pi_0}}$ and is a semi-abelian scheme whose special fiber modulo its maximum torus is a product of the Jacobians of I and I' respectively.

3.4. **Pulling back to arbitrary R_π.** Consider an arbitrary number field F as in section 2, containing $F_0 = \mathbb{Q}(\zeta_p)$, and let π be as in section 2. We will allow ourselves to enlarge F and change π repeatedly in what follows. Denote by

$$X_{R_\pi}, X_!, \mathcal{X}_!, A_{R_\pi}, A^0_{R_\pi}, \mathcal{A} \tag{3.1}$$

the objects obtained from the corresponding objects over R_{π_0} by base change via $R_{\pi_0} \subset R_\pi$. Note that the latter extension of rings can be ramified so A_{R_π} is not necessarily the Néron model of A_{K_π}; but all the schemes (respectively formal schemes) above continue to be smooth over R_π and $A^0_{R_\pi}$ is semiabelian, equal to $Pic^0_{X_1(Np)/R_\pi}$.

3.5. **Schemes attached to a fixed complex newform.** Let $f \in S_2(\Gamma_1(Np), \mathbb{C})$ be a newform and assume we are given an embedding $\rho : \mathcal{O}_f \to R_\pi$ with π as above. Let $A_{K_\pi} \to A_{f/K_\pi}$ be the induced morphism of abelian varieties over K_π obtained by base change from the morphism of abelian varieties over $\mathbb{Q}$, $J_1(Np)_{\mathbb{Q}} \to A_{f/\mathbb{Q}}$. Let A_{f/R_π} be the Néron model of A_{f/K_π}, let A^0_{f/R_π} be the connected component of the latter and let $A^0_{R_\pi} \to A^0_{f/R_\pi}$ be the induced morphism. By the theory of Raynaud extensions [13], pp. 33-34, there exists a group scheme C over R_π, which is an extension of an abelian scheme B over R_π by a torus T over R_π, such that $(A^0_{f/R_\pi})^\wedge \simeq \widehat{C}$. So we have an induced homomorphsim $\mathcal{A} \to (A^0_{f/R_\pi})^\wedge \to \widehat{B}$, hence an induced morphism $\mathcal{X}_! \to \widehat{B}$, hence an induced morphism $J^n_\pi(\mathcal{X}_!) \to J^n_\pi(B)$. Furthermore note that by the Néron property over R_{π_0}, plus base change from R_{π_0} to R_π, one has induced endomorphisms $T(l)_*$ of $A_{R_\pi}, A^0_{R_\pi}$, and $\mathcal{A}$ over R_π for all primes l. Similarly, by the Néron property over R_π, $\mathcal{O}_f$ acts on A_{f/R_π} and A^0_{f/R_π}. Clearly this action is compatible with the Hecke action on $A^0_{R_\pi}$. Similarly, for $d \in (\mathbb{Z}/p\mathbb{Z})^\times$, the diamond operators act on $\mathcal{X}_!$ and $\mathcal{A}$. By the functoriality of the Raynaud extension (cf. [13], pp. 33-34) we have an induced action of $\mathcal{O}_f$ on C and hence on B; so we have a ring homomorphism $\iota : \mathcal{O}_f \to End(B/R_\pi)$.

Assume first $B \neq 0$. Then, for any prime l, and any $d \in (\mathbb{Z}/p\mathbb{Z})^\times$, we have commutative diagrams

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{T(l)_*} & \mathcal{A} \\ \downarrow & & \downarrow \\ \widehat{B} & \xrightarrow{\iota(a_l)} & \widehat{B} \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathcal{X}_! & \xrightarrow{\langle d \rangle_p} & \mathcal{X}_! \\ \downarrow & & \downarrow \\ \mathcal{A} & \xrightarrow{\langle d \rangle_p} & \mathcal{A} \\ \downarrow & & \downarrow \\ \widehat{B} & \xrightarrow{\iota(\epsilon_p(d))} & \widehat{B} \end{array} \tag{3.2}$$

On the other hand assume $B = 0$; then $C = T$. So $\mathcal{O}_f$ acts on T via a homomorphism $\iota^\perp : \mathcal{O}_f \to End(T/R_\pi)$. Enlarging R_π again and replacing the schemes (3.1) by their pull-backs to this new R_π we may (and will!) assume T_{K_π} is split over K_π. So the identity component of its locally (!) finite type Néron model has the form $D = \mathbb{G}^\nu_{m/R_\pi}$ and, by the universality property of the Néron model, we get a homomorphism $T \to D$ over R_π. By the Néron property over R_π we have a homomorphism $\iota^\perp : \mathcal{O}_f \to End(D/R_\pi)$ and corresponding diagrams:

$$\begin{array}{ccc} \mathcal{A} & \xrightarrow{T(l)_*} & \mathcal{A} \\ \downarrow & & \downarrow \\ \widehat{D} & \xrightarrow{\iota^\perp(a_l)} & \widehat{D} \end{array} \quad \text{and} \quad \begin{array}{ccc} \mathcal{X}_! & \xrightarrow{\langle d \rangle_p} & \mathcal{X}_! \\ \downarrow & & \downarrow \\ \mathcal{A} & \xrightarrow{\langle d \rangle_p} & \mathcal{A} \\ \downarrow & & \downarrow \\ \widehat{D} & \xrightarrow{\iota^\perp(\epsilon_p(d))} & \widehat{D} \end{array} \tag{3.3}$$

3.6. **Lift of Igusa curve to characteristic zero** [10]**.** Recall that L denotes the line bundle on $X_1(N)_{R_p}$ such that the sections of the powers of L identify with the modular forms of various weights on $\Gamma_1(N)$; cf. [14] p. 450. Let $E_{p-1} \in H^0(X_1(N)_{R_p}, L^{p-1})$ be the normalized Eisenstein form of weight $p-1$ and let (ss) be the supersingular locus on $X_1(N)_{R_p}$, defined as the locus of E_{p-1}. Here E_{p-1} is normalized by the condition that its Fourier expansion has constant term 1. Take

an open covering

$$X_1(N) = \bigcup X_1(N)_i \tag{3.4}$$

such that L is trivial on each $X_1(N)_i$ and let x_i be a basis of L on $X_1(N)_i$. Then $E_{p-1} = \varphi_i x_i^{p-1}$ where $\varphi_i \in \mathcal{O}(X_1(N)_i)$. Set $x_i = u_{ij}x_j$, $u_{ij} \in \mathcal{O}^\times(X_1(N)_{ij})$, $X_1(N)_{ij} = X_1(N)_i \cap X_1(N)_j$. Set $z_i = x_i^{-1}$ (in [10] this z_i was denoted by t_i) and consider the R_π-scheme $X_1(N)_{!!}$ obtained by gluing the schemes

$$X_1(N)_{!!i} := Spec\ \mathcal{O}(X_1(N)_{i,R_\pi})[z_i]/(z_i^{p-1} - \varphi_i)$$

via $z_i = u_{ij}^{-1} z_j$ where

$$X_1(N)_{i,R_\pi} := X_1(N)_i \otimes_{R_p} R_\pi.$$

In the discussion below denote by an upper bar the functor $\otimes k$. Then $\overline{X_1(N)_{!!}}$ is clearly birationally isomorphic (hence isomorphic) to the Igusa curve I (cf. [14], pp. 460, 461). Now let $X \subset X_1(N)$ over R_p be an affine open set disjoint from the supersingular locus. Consider the open covering

$$X = \bigcup X_i \tag{3.5}$$

where $X_i = X_1(N)_i \cap X$. Consider the R_π-scheme $X_{!!} \subset X_1(N)_{!!}$ obtained by gluing the schemes $X_{!!i} := Spec\ \mathcal{O}(X_{i,R_\pi})[z_i]/(z_i^{p-1} - \varphi_i)$ via $z_i = u_{ij}^{-1} z_j$ (where $X_{i,R_\pi} := X_i \otimes_{R_p} R_\pi$). Note that $z_i^{p-1} - \varphi_i$ are monic polynomials whose derivatives are invertible in $\mathcal{O}(X_{i,R_\pi})[z_i]/(z_i^{p-1} - \varphi_i)$ so $X_{i!!}$ is an étale cover of X_{i,R_π} and is the pull back of the latter in $X_1(N)_{i!!}$. Note that the scheme $\overline{X_{!!}} = X_{!!} \otimes k$ is isomorphic to $\overline{\mathcal{X}}_! = \mathcal{X}_! \otimes k$. Recall from [10] that the isomorphism $\overline{X}_{!!} \simeq \overline{\mathcal{X}}_!$ lifts uniquely to an isomorphism $(X_{!!})^\wedge \simeq \mathcal{X}_!$. This follows immediately by applying the standard Lemma 3.2 below to $S := \mathcal{O}(\mathcal{X}_i)$, $\mathcal{X}_i = \widehat{X}_i$, $T = \mathcal{O}(\mathcal{X}_{!i})$, $\mathcal{X}_{!i} = e^{-1}(\mathcal{X}_i)$. □

Lemma 3.2. *Let $S \to T$ be a homomorphism of flat π-adically complete R_π-algebras, let $f \in S[z]$ be a monic polynomial and assume we have a homomorphism $\overline{\sigma} : \overline{S}[z]/(\overline{f}) \to \overline{T}$ such that df/dz is invertible in $\overline{S}[z]/(\overline{f})$. Then $\overline{\sigma}$ lifts uniquely to a homomorphism $\sigma : S[z]/(f) \to T$. If $\overline{\sigma}$ is an isomorphism then so is σ.*

Proof. One defines σ by sending the class $\xi \in S[z]/(f)$ of z into the unique solution $\tau \in T$ (which exists by Hensel) of the equation $f = 0$ with $\overline{\sigma}(\overline{\xi}) = \overline{\tau}$, where the upper bar means class mod π. □

3.7. **Serre lifts.** Let M be any positive integer. In what follows a *classical modular form* over a ring B, of weight κ, on $\Gamma_1(M)$ will be understood in the sense of [14] as a rule that attaches to any B-algebra C and any triple consisting of an elliptic curve E/C, an embedding $\mu_{M,C} \to E[M]$, and an invertible one form on E an element of C satisfying the usual compatibility rules and the usual holomorphy condition for the Fourier expansion (evaluation on the Tate curve). We denote by $M_\kappa(\Gamma_1(M), B)$ the B-module of all these forms and by $S_\kappa(\Gamma_1(M), B)$ the module of corresponding cusp forms. For the concepts of Hecke operators, diamond operators, and newforms used below we refer to [14]. Let $\Theta_p : (\mathbb{Z}/p\mathbb{Z})^\times \to \mathbb{Z}_p^\times \subset R_\pi^\times$ be the Teichmüller character. We have the following "reduction to weight 2" result [14], p. 478:

Proposition 3.3. *Let $(N, p) = 1, N > 4$ and let $\overline{f} \in S_\kappa(\Gamma_1(N), k)$ be a newform, where $3 \leq \kappa \leq p$. Then there exists a newform $f \in S_2(\Gamma_1(Np), \mathbb{C})$ with character ϵ and there exists an embedding $\rho : \mathcal{O}_f \to R_\pi$ such that:*

1) The image of f via $\mathcal{O}_f \xrightarrow{\rho} R_\pi \to k$ is $\overline{f}$;
2) $\rho \circ \epsilon_p = \Theta_p^{\kappa-2}$.

In the above ϵ_p is the "p-part" in the decomposition $\epsilon = \epsilon_p \epsilon_N$. We shall refer to the pair (f, ρ), or simply to f, as a *Serre lift* of $\overline{f}$. Recall from [14] that Serre lifts are not unique.

4. Construction of differential modular forms

4.1. **δ_π-characters.** Let B be an abelian scheme over R_π. Recall that by a δ_π-character of B, of order n, we understand a homomorphism $J^n_\pi(B) \to \widehat{\mathbb{G}_a}$.

Proposition 4.1. *Consider an abelian scheme B over R_π of relative dimension g. Then the R_π-module of δ_π-characters of B of order 2 has rank $\geq g$.*

In [3] this was proved under the assumption that the ramification index of R_π satisfies $e \leq p-2$; the method there does not seem to extend to arbitrary e. The argument below, for general e, follows a method in [5].

Proof. Let $\mathcal{F} \in R_\pi[[T_1, T_2]]^g$ be the formal group law of B and $l(T) \in K_\pi[[T]]^g$ its logarithm; here T, T_1, T_2 are g-tuples of variables. For any $n \geq 1$ let

$$L^n_\pi = \frac{1}{\pi} l(\phi^n(T))_{|T=0} \in K_\pi[[T, \delta_\pi T, ..., \delta^n_\pi T]]$$

where $\phi : K_\pi[[T, \delta_\pi T, ...]] \to K_\pi[[T, \delta_\pi T, ...]]$,

$$\phi(T) = T^p + \pi \delta_\pi T, \ \ \phi(\delta_\pi T) = (\delta_\pi T)^p + \pi \delta^2_\pi T, ...$$

We claim that there is an integer $\nu \geq 1$ such that

$$\pi^\nu L^n_\pi \in R_\pi[[T]][\delta_\pi T, ..., \delta^n_\pi]\hat{}\,;$$

and, by the way, if in addition $e \leq p-1$ then one can take $\nu = 0$. Indeed by [15], p. 64, we have

$$l(T) = \sum_{|\alpha| \geq 1} \frac{A_\alpha}{|\alpha|} T^\alpha$$

for some $A_\alpha \in R_\pi$. Hence setting $G^n_\pi = \frac{1}{\pi} \phi^n(T)_{|T=0}$ we have

$$L^n_\pi = \frac{1}{\pi} \sum_{|\alpha| \geq 1} \phi^n(A_\alpha) \frac{\pi^{|\alpha|-1}}{|\alpha|} (G^n_\pi)^\alpha$$

Now if v_π is the π-adic valuation we have

$$v_\pi \left(\frac{\pi^{|\alpha|-1}}{|\alpha|} \right) \geq |\alpha| - 1 - e \cdot \log_p |\alpha|$$

So the above goes to ∞ for $|\alpha| \to \infty$; moreover the above is ≥ 0 if $e \leq p-1$. This proves the claim. Let N^2 be the kernel of $J^2_\pi(B) \to \widehat{B}$. Exactly as in [5] the g-tuples $\pi^\nu L^2_\pi$ and $\pi^\nu L^1_\pi$ define $2g$ homomorphisms $N^2 \to \widehat{\mathbb{G}_a}$ that are R_π-linearly independent. So $Hom(N^2, \widehat{\mathbb{G}_a})$ has rank $\geq 2g$. We are done by considering the standard exact sequence (cf. [3]):

$$0 = H^0(B, \mathcal{O}) \to Hom(J^2_\pi(B), \widehat{\mathbb{G}_a}) \to Hom(N^2, \widehat{\mathbb{G}_a}) \to H^1(B, \mathcal{O}) \simeq R^g_\pi.$$

□

Remark 4.2. Recall from [3] that $\mathbb{G}_m = Spec R_\pi[x, x^{-1}]$ has a remarkable δ_π-character, i.e. a homomorphism $\psi : J^1_\pi(\mathbb{G}_m) \to \widehat{\mathbb{G}_a}$ defined by

$$\psi = \pi^m \sum_{n \geq 1} (-1)^{n-1} \frac{\pi^n}{n} \left(\frac{\delta_\pi x}{x^p} \right)^n$$

for an appropriate $m \in \mathbb{Z}$. So if $D = \mathbb{G}_a^\nu$ then there is a canonical system $\psi_1, ..., \psi_\nu : J^1_\pi(D) \to \widehat{\mathbb{G}_a}$ of δ_π-characters obtained by composing ψ above with the canonical projections onto the factors.

4.2. **Semi-invariant δ_π-characters.** Assume the notation of Section 3.5 and let $\rho : K_f(\zeta_N, \zeta_p) \to K_\pi$ be an embedding. (One can always choose π such that such a ρ exists.) So $\rho(\mathcal{O}_f[1/N, \zeta_N, \zeta_p]) \subset R_\pi$. Then, exactly as in [7], Proposition 4.5 and using the same argument as on the top of page 992, loc. cit., we get:

Lemma 4.3. *Assume $B \neq 0$ and let $\mathcal{T} = \iota(\mathcal{O}_f) \subset End(B/R_\pi)$. Then there exists a non-zero δ_π-character $\psi \in \mathcal{O}(J^2_\pi(B))$ and a ring homomorphism $\chi : \mathcal{T} \to R_\pi$ such that $\psi \circ \tau = \chi(\tau) \cdot \psi$ for all $\tau \in \mathcal{T}$.*

Similarly we get:

Lemma 4.4. *Assume $B = 0$ (hence D is defined and non-zero). Let $\mathcal{T} = \iota^\perp(\mathcal{O}_f) \subset End(D/R_\pi)$. Then there exists a non-zero δ_π-character $\psi \in \mathcal{O}(J^1_\pi(D))$ and a ring homomorphism $\chi : \mathcal{T} \to R_\pi$ such that $\psi \circ \tau = \chi(\tau) \cdot \psi$ for all $\tau \in \mathcal{T}$.*

4.3. **δ_π-modular forms from $\mathcal{X}_!$.** Recall from [10] the following:

Proposition 4.5. *There is a natural isomorphism*

$$\mathcal{O}(J^r_\pi(\mathcal{X}_!)) \simeq \bigoplus_{\kappa=0}^{p-2} M^r_\pi(-\kappa).$$

Via this isomorphism $M^r_\pi(-\kappa)$ is identified with the subspace of all $\varphi \in \mathcal{O}(J^r_\pi(\mathcal{X}_!))$ on which the diamond operators $\langle d \rangle_p$ with $d \in (\mathbb{Z}/p\mathbb{Z})^\times$ act by the character $d \mapsto \Theta_p(d)^{-\kappa}$ (i.e. $\langle d \rangle_p \varphi = \Theta_p(d)^{-\kappa} \varphi$.)

The ring $\mathcal{O}(J^n_\pi(\mathcal{X}_!))$ can be referred to as the ring of *δ_π-modular forms* of level Np, order n, and weight 0.

Let now f be as in section 3.5 and let $\rho : \mathcal{O}_f[\zeta_{Np}] \to R_\pi$ be an embedding. Assume the notation in section 3.5 and let $\epsilon = \epsilon_N \epsilon_p$ be the character of f.

Assume first $B \neq 0$.

Let $\psi : J^2_\pi(B) \to \widehat{\mathbb{G}_a} = \widehat{\mathbb{A}^1}$ and χ be as in Lemma 4.3. So for $d \in (\mathbb{Z}/(p\mathbb{Z})^\times$ we have

$$\psi \circ \iota(\epsilon_p(l)) = \chi(\iota(\epsilon_p(d))) \cdot \psi.$$

Then if $\beta : \mathcal{X}_! \to \widehat{B}$ is the vertical morphism in equation 3.2 and if $\beta^2 : J^2_\pi(\mathcal{X}_!) \to J^2_\pi(\widehat{B}) = J^2_\pi(B)$ is the induced morphism set

$$f^\sharp = \psi \circ \beta^2 : J^2_\pi(\mathcal{X}_!) \to \widehat{\mathbb{A}^1} \tag{4.1}$$

We get

$$\langle d \rangle_p f^\sharp = \psi \circ \beta^2 \circ \langle d \rangle_p = \psi \circ \iota(\epsilon_p(d)) \circ \beta^2 = \chi(\iota(\epsilon_p(d))) \cdot \psi \circ \beta^2 = \chi(\iota(\epsilon_p(d))) \cdot f^\sharp.$$

Set $\rho' := \chi \circ \iota : \mathcal{O}_f \to R_\pi$. Then $f^\sharp$ has weight $-\kappa'$ for some integer $0 \leq \kappa' \leq p-2$ if and only if

$$\rho' \circ \epsilon_p = \Theta_p^{-\kappa'}.$$

If in addition f and ρ are as in Proposition 3.3, so in particular $\overline{f}$ has weight κ, then

$$\rho \circ \epsilon_p = \Theta_p^{\kappa-2}.$$

Let $g \in (\mathbb{Z}/p\mathbb{Z})^\times$ be a generator; so $\Theta_p(g) \in \mathbb{Z}_p^\times$ is a primitive $(p-1)$-th root of unity in $\mathbb{Q}_p$ and $\epsilon_p(g)$ is a (not necessarily primitive) $(p-1)$-th root of unity in $\mathbb{C}$. Then ρ, ρ' induce embeddings $r, r' : \mathbb{Q}(\epsilon_p(g)) \to \mathbb{Q}(\Theta_p(g))$ hence $r(\epsilon_p(g)) = \Theta_p(g)^{\kappa-2}$. Now since $\mathbb{Q}(\Theta_p(g))/\mathbb{Q}$ is Galois there is an automorphism γ of this extension such that $\gamma \circ r = r'$; by the structure of the Galois group we must have $\gamma(\Theta_p(g)) = \Theta_p(g)^c$ for some integer c coprime to $p-1$. We get

$$\rho'(\epsilon_p(g)) = r'(\epsilon_p(g))) = \gamma(r(\epsilon_p(g)) = \Theta_p(g)^{c(\kappa-2)}$$

hence $\rho' \circ \epsilon_p = \Theta_p^{c(\kappa-2)} = \Theta_p^{-\kappa'}$, where κ' is the unique integer between 0 and $p-2$ such that $\kappa' \equiv c(2-\kappa) \bmod p-1$. Then we have proved the following:

Proposition 4.6. *Let $\overline{f} \in S_\kappa(\Gamma_1(N), k)$ be a newform as in Proposition 3.3 and let (f, ρ) be a Serre lift (cf. loc.cit.). Assume $B \neq 0$ and let $f^\sharp$ be as in 4.1. Then $f^\sharp \in M_\pi^2(-\kappa')$ where κ' is a conjugate of κ.*

Similarly if $B = 0$ (hence D is defined and non-zero) we let $\psi : J_\pi^1(D) \to \widehat{\mathbb{G}_a} = \widehat{\mathbb{A}^1}$ and χ be as in Lemma 4.4. Then if $\beta : \mathcal{X}_! \to \widehat{D}$ is the vertical morphism in equation 3.3 and if $\beta^1 : J_\pi^1(\mathcal{X}_!) \to J^2(\widehat{D}) = J_\pi^2(D)$ is the induced morphism we set

$$f^\sharp = \psi \circ \beta^1 : J_\pi^1(\mathcal{X}_!) \to \widehat{\mathbb{A}^1} \tag{4.2}$$

and we get:

Proposition 4.7. *Let $\overline{f} \in S_\kappa(\Gamma_1(N), k)$ be a newform as in Proposition 3.3 and let (f, ρ) be a Serre lift (cf. loc.cit.). Assume $B = 0$ and let $f^\sharp$ be as in 4.2. Then $f^\sharp \in M_\pi^1(-\kappa')$ where κ' is a conjugate of κ.*

The proof of Theorem 1.1 will be concluded if we prove the following Proposition, where we assume that the embedding $\rho : \mathcal{O}_f \to R_\pi$ is extended (which is always possible by changing π) to an embedding of the Galois closure of K_f into K_π.

Proposition 4.8. *Let $f^\sharp$ be as in 4.1 or 4.2. Then $f^\sharp$ is non-zero and its δ_π-Fourier expansion has the form*

$$E(f^\sharp) = \sum_\sigma P_\sigma(\phi)(f^{(-1)}(q))^\sigma,$$

with $P_\sigma(\phi)$ polynomials in $K_\pi[\phi]$ and σ running through the set of all embeddings of K_f into $\mathbb{C}$.

Proof. We have $f^\sharp \neq 0$ because $\psi \neq 0$ and the images of $\beta : \mathcal{X}_! \to \widehat{B}$ and $\beta : \mathcal{X}_! \to \widehat{D}$ generate $\widehat{B}$ and $\widehat{D}$ respectively. The shape of $E(f^\sharp)$ follows directly from the way $f^\sharp$ was defined, exactly as in [7], proof of Theorem 6.3. For convenience we sketch here the argument. Assume $B \neq 0$; the case $B = 0$ is similar (with D playing the role of B). Let g be the dimension of B, let $\omega_1, ..., \omega_g$ be a basis of global 1-forms on B and consider the formal logarithm $L = L(T_1, ..., T_g)$ of B

with components $L_1, ..., L_g$ (which are series in the variables $T_1, ..., T_g$) such that $\omega_i = dL_i$. One can write

$$\beta^* \omega_i = \sum_\sigma c_{i\sigma} \sum_n a_n^\sigma q^{n-1} dq,$$

with $c_{i\sigma} \in R_\pi$. On the other hand, if $\beta^*(T_i) = \varphi_i(q) \in R_\pi[[q]]$ and $\varphi = (\varphi_1, ..., \varphi_g)$, we have

$$\beta^* \omega_i = \beta^* \left(\sum_j \frac{dL_i}{dT_j} dT_j \right) = \sum_j \frac{dL_i}{dT_j}(\varphi(q)) \frac{d\varphi_j}{dq} dq = \frac{d}{dq}(L_i(\varphi(q))dq.$$

One then gets

$$L_i(\varphi(q)) = \sum_\sigma c_{i\sigma} \sum_n a_n^\sigma \frac{q^n}{n} = \sum_\sigma c_{i\sigma} (f^{(-1)}(q))^\sigma.$$

On the other hand one has

$$\psi = \sum_i Q_i(\phi) L_i$$

for some polynomials $Q_i(\phi) \in K_\pi[\phi]$. We get

$$f^\sharp = \beta^* \psi = \sum_i Q_i(\phi) \left(\sum_\sigma c_{i\sigma} (f^{(-1)}(q))^\sigma \right) = \sum_\sigma \left(\sum_i c_{i\sigma} Q_i(\phi) \right) (f^{(-1)}(q))^\sigma,$$

and we are done by setting $P_\sigma(\phi) = \sum_i c_{i\sigma} Q_i(\phi)$. □

5. Link with Igusa differential modular functions

The aim of this section is to construct a natural (and somewhat unexpected) ring homomorphism from the ring $\mathcal{O}(J_\pi^n(\mathcal{X}_!))$ of δ_π-modular forms of level Np, order n, and weight 0 to the ring $S_\heartsuit^{n+1} \otimes_{R_p} R_\pi$ where $S_\heartsuit^{n+1}$ is the ring of *Igusa δ_p-modular forms of level N* [9]. We start by reviewing the rings $S_\heartsuit^n$.

Recall our basic setting in section 2. With $X \subset X_1(N)$ disjoint from the supersingular locus, and with L and $V \to X$ as in section 2 set:

$$S^n = \mathcal{O}(J_p^n(X)),\ M^n = \mathcal{O}(J_p^n(V)),\ S^\infty = \lim_{\to} S^n,\ M^\infty = \lim_{\to} M^n;$$

the elements of M^n are the δ_p-modular functions of order n. For $w \in W$ recall that we let $M^n(w)$ be the the R_p-module of all $f \in M^n$ such that $\lambda \star f = \lambda^w f$ for $\lambda \in R_p^\times$, where $\star$ is the natural action of the multiplicative group; the elements of $M^n(w)$ are the δ_p-modular forms of weight w. On the other hand recall the natural δ_p-expansion map

$$E : M^n \to R_p((q))^n := R_p((q))[q', ..., q^{(n)}]\hat{};$$

its restriction to each $M^n(w)$ is injective. Let $S_\heartsuit^n \subset R_p((q))^n$ be the image of M^n via E and let

$$S_\heartsuit^\infty = \lim_{\to} S_\heartsuit^n \subset R_p((q))^\infty := \lim_{\to} R_p((q))^n.$$

The ring $S_\heartsuit^\infty$ can be referred to as the ring of *Igusa δ_p-modular functions of level N* (or the ring of the δ_p-Igusa curve of level N [9]). Recall from [1] and [6], p. 269, that there is a (necessarily unique) $f^\partial \in M^1(\phi - 1)$ such that $E(f^\partial) = 1$. Moreover we have a congruence

$$f^\partial \equiv E_{p-1} \mod p \text{ in } M^1. \tag{5.1}$$

Theorem 5.1. *There exists a sequence of S^0-algebra homomorphisms,*

$$\mathcal{O}(J^n_\pi(\mathcal{X}_!)) \to S^{n+1}_\heartsuit \otimes_{R_p} R_\pi,$$

compatible with each other as n varies, and commuting with δ_π.

As we shall see these homomorphisms are "natural". Also if we consider the limit

$$\mathcal{O}(J^\infty_\pi(\mathcal{X}_!)) := \lim_{\overrightarrow{n}} \mathcal{O}(J^n_\pi(\mathcal{X}_!))$$

our Theorem 5.1 yields an S^0-algebra homomorphism, commuting with δ_π,

$$\mathcal{O}(J^\infty_\pi(\mathcal{X}_!)) \to S^\infty_\heartsuit \otimes_{R_p} R_\pi. \tag{5.2}$$

Proof of Theorem 5.1. We begin by constructing an S^0-algebra homomorphism

$$\mathcal{O}(\mathcal{X}_!) \to S^1_\heartsuit \otimes_{R_p} R_\pi. \tag{5.3}$$

Indeed consider a cover as in Equation 3.5 and, using the notation preceding that equation, set $f^\partial = \Phi_i x_i^{\phi-1}$ where $\Phi_i \in S^1_i := \mathcal{O}(J^n_p(X_i))$. Let

$$M^1_i = S^1_i[x_i, x_i^{-1}, \delta_p x_i]^\wedge = S^1_i[z_i, z_i^{-1}, \delta_p z_i]^\wedge$$

Then

$$M^1_i/(f^\partial - 1) \simeq S^1_i[z_i, z_i^{-1}, \delta_p z_i]^\wedge/(z_i^{\phi-1} - \Phi_i).$$

Since the latter is Noetherian and p-adically complete, and since

$$(M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi$$

is finite over it, it follows that $(M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi$ is p-adically complete, hence π-adically complete. Recall that, setting $S_i = \mathcal{O}(X_i)$, we have

$$\mathcal{O}(\mathcal{X}_{!i}) = \frac{(S_i \otimes_{R_p} R_\pi)[z_i]^\wedge}{(z_i^{p-1} - \varphi_i)}, \quad (M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi = \frac{S^1_i[z_i, z_i^{-1}, \delta_p z_i]^\wedge \otimes_{R_p} R_\pi}{(z_i^{\phi-1} - \Phi_i)}.$$

Hence

$$\overline{\mathcal{O}(\mathcal{X}_{!i})} = \frac{\overline{S_i}[z_i]}{(z_i^{p-1} - \overline{\varphi_i})}, \quad \overline{(M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi} = \frac{\overline{S^1_i}[z_i, z_i^{-1}, \delta_p z_i]}{(z_i^{p-1} - \overline{\Phi_i})}.$$

Since, by Equation 5.1, $\overline{\Phi_i} = \overline{\varphi_i}$ in $\overline{S^1_i}$ and $\overline{S_i} \subset \overline{S^1_i}$ we get natural inclusions, compatible with varying i:

$$\overline{\mathcal{O}(\mathcal{X}_{!i})} \subset \overline{(M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi}$$

and hence, by Lemma 3.2, unique liftings of the above to homomorphisms

$$\mathcal{O}(\mathcal{X}_{!i}) \to (M^1_i/(f^\partial - 1)) \otimes_{R_p} R_\pi.$$

By uniqueness these liftings glue together to give a homomorphism

$$\mathcal{O}(\mathcal{X}_!) \to (M^1/(f^\partial - 1)) \otimes_{R_p} R_\pi.$$

Composing with the natural map

$$(M^1/(f^\partial - 1)) \otimes_{R_p} R_\pi \to S^1_\heartsuit \otimes_{R_p} R_\pi$$

we get the desired homomorphism 5.3

Now by the universality property of $(\mathcal{O}(J^n_\pi(\mathcal{X}_!)))_{n\geq 0}$ and by the fact that

$$(S^{n+1}_\heartsuit \otimes_{R_p} R_\pi)_{n\geq 0}$$

is a δ_π-prolongation sequence we get that the homomorphism 5.3 induces natural compatible ring homomorphisms commuting with δ_π as in the statement of the theorem. □

Remark 5.2. It would be interesting to see if the homomorphism 5.2 is injective. On the other hand we claim that its reduction mod π,

$$\overline{\mathcal{O}(J^\infty_\pi(\mathcal{X}_!))} \to \overline{S^\infty_\heartsuit}, \tag{5.4}$$

is not injective. Indeed, since $X_{!!}$ is étale over X it follows from [3], Proposition 1.4, that

$$\mathcal{O}(J^n_\pi(\mathcal{X}_{!i})) = \mathcal{O}(\mathcal{X}_{!i})[\delta_\pi z_i, ..., \delta^n_\pi z_i]\hat{}$$

hence

$$\overline{\mathcal{O}(J^n_\pi(\mathcal{X}_{!i}))} = \overline{\mathcal{O}(\mathcal{X}_{!i})}[\delta_\pi z_i, ..., \delta^n_\pi z_i].$$

On the other hand by [9], Introduction, we have

$$\overline{S^\infty_\heartsuit \otimes_{R_p} R_\pi} = \overline{S^\infty_\heartsuit} = \frac{\overline{S^\infty}[z_i, \delta_p z_i, \delta^2_p z_i, ...]}{(f^\partial - 1, \delta_p(f^\partial - 1), \delta^2_p(f^\partial - 1), ...)}.$$

So the map 5.4 sends $\delta^a_\pi z_i$ into 0 for all $1 \leq a \leq p-2$; in particular the map 5.4 is not injective.

References

1. Barcau, M: Isogeny covariant differential modular forms and the space of elliptic curves up to isogeny, Compositio Math. 137, 237-273 (2003)
2. Bosch S., Lütkebohmert W., Raynaud M., Néron models, Springer 1990.
3. Buium, A.: Differential characters of Abelian varieties over $p-$adic fields, Invent. Math. 122, 309-340 (1995).
4. Buium, A.: Differential modular forms, Crelle J., 520 (2000), 95-167.
5. Buium, A.: Geometry of Fermat adeles, Trans. AMS 357, 3 (2005), 901-964.
6. Buium, A.: Arithmetic Differential Equations. Math. Surveys and Monographs 118, AMS (2005)
7. Buium A., Differential eigenforms, J. Number Theory 128 (2008), 979-1010.
8. Buium, A., Poonen, B., Independence of points on elliptic curves arising from special points on modular and Shimura curves, II: local results, Compositio Math., 145 (2009), 566-602.
9. Buium A., Saha A., The ring of differential Fourier expansions, J. of Number Theory 132 (2012), 896-937.
10. Buium, A., Saha, A., Differential overconvergence, in: Algebraic methods in dynamical systems; volume dedicated to Michael Singer's 60th birthday, Banach Center Publications, Vol 94, 99-129 (2011).
11. Buium, A., Poonen, B.: Independence of points on elliptic curves arising from special points on modular and Shimura curves, II: local results, Compositio Math., 145 (2009), 566-602.
12. Diamond, F., and Im, J.: Modular forms and modular curves. In:Seminar on Fermat's Last Theorem, Conference Proceedings, Volume 17, Canadian Mathematical Society, pp. 39-134 (1995).
13. Faltings, G., Chai, C-L., Degeneration of Abelian Varieties, Springer 1990.
14. Gross, B. H., A tameness criterion for Galois representations associated to modular forms mod p, Duke Math. J., 61, 2, 445-517 (1990)
15. Hazewinkel, Formal Groups and Applications, AMS, 2012.

University of New Mexico, Albuquerque, NM 87131
E-mail address: buium@math.unm.edu, arnab@math.unm.edu